


10th IAPC Meeting

Tenth World Conference on
Physico-Chemical Methods in Drug Discovery
&
Sixth World Conference on ADMET and DMPK



Book of Abstracts



September 2023 :: Belgrade, Serbia

10th IAPC Meeting

*Tenth World Conference on Physico-Chemical Methods in Drug Discovery
&*

*Sixth World Conference on ADMET and DMPK
Belgrade, Serbia, September 4-6, 2023*

Book of Abstracts

Organized by

International Association of Physical Chemists

&

Faculty of Chemistry, University of Belgrade, Serbia

Published by

International Association of Physical Chemists

E-mail: office@iapchem.org, URL: <http://www.iapchem.org>

For Publisher

Zoran Mandić

Editor

Tatjana Verbić & Zoran Mandić

Design, page making and computer layout

Aleksandar Dekanski

On Line version only

The Scientific and Organizing Committee:

Tatjana Verbić, Conference Chair
University of Belgrade, Serbia,

Alex Avdeef
ADME Research, New York, USA

Kiyohiko Sugano
Ritsumeikan University, Osaka, Japan

Kin Tam
University of Macau, Macau

Zoran Mandić
University of Zagreb, Croatia

Klara Valko
Biomimetic chromatography Ltd. UK

Godefridus J. Peters
Amsterdam University Medical Centers, The Netherlands

Hong Wan
WHDeX Consulting AB, Sweden

Local Organizing Committee

Tatjana Verbić, Conference Chair
University of Belgrade, Faculty of Chemistry, Serbia

Goran Roglić
University of Belgrade, Faculty of Chemistry, Serbia

Ilija Cvijetić
University of Belgrade, Faculty of Chemistry, Serbia

Miloš Pešić
University of Belgrade, Faculty of Chemistry, Serbia

Olivera Marković
University of Belgrade, Institute of Chemistry, Technology and Metallurgy, Serbia

Aleksandar Dekanski
University of Belgrade, Institute of Chemistry, Technology and Metallurgy, Serbia

Marija Popović Nikolić
University of Belgrade, Faculty of Pharmacy, Serbia

Organization of IAPC-10 Meeting is supported by



Ministry of Science, Technological
Development and Innovation
of the Republic of Serbia

Sponsors and Exhibitors



Nenad M. Kostić
Fund for Chemical Sciences



Granny's
Secret

Bakina
Tajna



SUPERLAB[®]
Your lab - Our passion



ADMET



CONTENTS

Oral presentations

Virtual assessment to provide insights into drug-exciipient-intestinal fluid interactions and support virtual formulation design <i>Christel Bergström</i>	2
In vitro evaluation of drug presence in the micellar phase of contents of upper small intestine: Rationale, challenges, opportunities <i>Christos Reppas</i>	3
Continuous improvement in the molecular properties assessment world <i>Gilles Goetz, Jim Federico, Brendon Kapinos, Matt Troutman</i>	4
Nanotechnology enabled innovation in inhalation drug delivery <i>Eride Quarta, Paolo Colombo, Lorenzo Degli Esposti, Daniele Catalucci, Michele Iafisco, Claudio De Luca, Gaia Colombo</i>	5
Neuro pharmacokinetics: the secret life of - old and novel - psychopharmacological drugs <i>Miroslav Savić</i>	6
Celebrating 40 years of the ROSS™ electrode and 50 years of pH_{max} <i>Alex Avdeef</i>	7
Salt disproportionation in bicarbonate buffer <i>Kiyohiko Sugano</i>	8
Development of rapidly dissolving 3D-printed tablets for personalized medicine by applying acid-base supersolubilization (ABS) principle <i>Abu Serajuddin</i>	9
Salt selection for development - solubility & bioperformance advantages from salts made by weaker acidic counterions explained by thermodynamic equilibria <i>Yan He</i>	10
Solubility prediction by deep learning of quantum information <i>Tonglei Li</i>	11
Micro-scale solubility measurement to guide early decision making <i>Sami Svanbäck</i>	12
Insights in the solubility and dissolution rate determination of cocrystals <i>Elisabet Fuguet, Hanan Fael, Rafael Barbas, Rafel Prohens, Rebeca Ruiz, Clara Ràfols</i>	13

Preparation and characterization of solid dispersion of mesalazine using anti-solvent precipitation technique

Emin, Zekir Kade, Poturcu, Alan Bogos, Aygme, Konstantyn Kozłowski, Abdalla E. Elbashir _____ 14

Revealing the story of an orphan drug: clofazimine speciation and solubilization as a function of pH

Tatjana Ž. Verbić, Alex Avdeef, Kin Y. Tam, Olivera S. Marković, Miloš P. Pešić, Igor A. Topalović, Dušan Ž. Veljković, Mufaddal Kathawala, Abu T. M. Serajuddin __ 15

Amorphous solid dispersions: True supersaturation measured in a time-resolved manner by microdialysis

Annette Bauer Brandl _____ 16

Metabolism as a target for drug development

Godefridus J. Peters, Kin Tam, Marika Franczak, Yan Zhou, Filippo Minutolo, Carlotta Granchi, R. Tom Smolenski, Elisa Giovannetti _____ 17

Dichloroacetophenone derivatives as pyruvate dehydrogenase kinase inhibitor in non-small cell lung cancer models

Kin Tam _____ 18

Development of catalytic inhibitors of topoisomerase II α as chemotherapeutic agents

Barbara Herlah, Matej Janežič, Kaja Bergant Loboda, Katja Valjavec, Andrej Perdih _ 19

Highlight on the benefits of PBPK modeling: A link between drug properties and its in vivo performance

Sandra Cvijić _____ 20

Navigating towards improved cytotoxicity assessment in nanomedicine development: Shifting from colorimetric to fluorescence-based assays

Ines Nikolić, Marija Petrović, Jelena Mitović, Emmanuelle Sublet, Olivier Jordan, Snežana Savić, Gerrit Borchard _____ 21

Chromatography platform to highlight compounds behavior in a membrane: A journey of interactions

Laurence Philippe-Venec _____ 22

Combining biomimetic chromatography and the quantitative structure-(chromatographic) retention relationships approach using machine learning

Krzysztof Ciura _____ 24

Application of lipophilicity, protein and phospholipid binding measured by HPLC for prediction of ADME properties and toxicity of compounds

Klara Valko, Bard Calvin _____ 25

Application of biomimetic chromatography for the prediction of acute aquatic toxicity of organic pollutants

C. Stergiopoulos, F. Tsopeas, M. Ochsenkühn-Petropoulou, K. Valko _____ 26

From $\Delta \log k_w^{IAM}$ to multidimensional biomimetic chromatography: a journey to shed light on the absorption potential of therapeutics. What will the future bring?	
<i>Giacomo Russo</i> _____	27
Ability of physicochemical systems to predict skin permeation of neutral compounds. A comparison study	
<i>Martí Rosés, Sara Soriano-Meseguer, Adriana Port, Elisabet Fuguet</i> _____	28
Towards better understanding of drug interactions with mesoporous silica carriers by using inverse gas chromatography and molecular modelling	
<i>Andreas Niederquell, Barbora Vraníková, Martin Kuentz</i> _____	29
In silico approaches for understanding the role of intermolecular interactions in formulations and combination therapies: implications for ADME properties optimization	
<i>Mire Zloh</i> _____	30
Octanol-water distribution coefficient ($\log D$) as molecular descriptor to count environmental effect in QSAR models: comparison on experimental and predicted values for common reference drug substances for wide pH scale	
<i>Mare Oja, Uko Maran</i> _____	31
With a little help from computer-aided drug design – new antitumor agents as tubulin polymerization inhibitors	
<i>Robert Vianello</i> _____	32
Elucidation of molecular mechanisms of activity of <i>Echinacea spp.</i> constituents for possible treatment of COVID 19 by computer-aided methods	
<i>Slavica Erić, Milena Jović, Mire Zloh</i> _____	33
Quantitative structure-activity relationship of pyrimidine and uracil derivatives for characterization and evaluation the potential of chemical compounds for cervical cancer	
<i>S. Zukic, U. Maran</i> _____	34
Bridging science & regulation: quality by design in patient- focused formulation development	
<i>Jelena Đuriš</i> _____	35
Improvement of digital twin for transdermal fentanyl delivery based on anomalous diffusion	
<i>Milena Čukić Radenković, Slobodanka Galović</i> _____	36
Sponsors' presentations	
High-resolution mass spectrometry in drug discovery and design – trends and perspectives	
<i>Luka Mihajlović</i> _____	38

Determination of active components in medicines and food on Shimadzu chromatographies	
<i>Stefan Jovanović, Aleksandar Zurkić, Vanja Vranjevac</i>	39
Advancing drug safety: PrimeVigilance's comprehensive pharmacovigilance approach	
<i>Nebojša Jankov, Nikola Stojanović</i>	40
Poster presentations	
Evaluation of the micellization of the biosurfactant sodium taurocholate using fluorescence measurements	
<i>S. Amézqueta, E. Fuguet, U. Casanova, C. Ràfols</i>	42
Investigation of molecular mechanism of action of <i>Satureja Montana</i> essential oil constituents related to effects on viremias	
<i>Slavica Erić, Đorđe Vasilić, Katarina Ilić, Mire Zloh</i>	43
Computational study of interactions of <i>Cannabis Sativa</i> constituents with potential epigenetic targets involved in processes of multiple sclerosis	
<i>Slavica Erić, Charline Hendricks, Mire Zloh</i>	44
Computational study of the monoamin oxidase B mechanism- based irreversible inhibitors	
<i>Lucija Vrban, Robert Vianello</i>	45
Virtual docking and design of novel Histone deacetylase and Rho- associated protein kinases dual inhibitors (HDAC/ROCKs)	
<i>Milan Beljkaš, Miloš Petković, Katarina Nikolić, Slavica Oljačić</i>	46
Towards the multitarget HDAC Inhibitors for the treatment of pancreatic carcinoma by joining the drug synergy predictions and the molecular modeling	
<i>Nemanja Đoković, Aleksandra Ilić, Alen Čebzan, Branko Radović, Dušan Ružić, Ana Đurić, Tatjana Srdić-Rajić, Katarina Nikolić</i>	47
Biomimetic characteristics of dual TLC retention mechanism	
<i>Darija Obradović, Lukasz Komsta, Marija Popović-Nikolić, Jovana Milutinović, Saša Lazović</i>	48
Study of ionization of montelukast in differently charged micellar solutions as biomembrane mimetic systems	
<i>Marija Popović-Nikolić, Slavica Oljačić, Katarina Nikolić, Gordana Popović</i>	49
Improvement of the vitamin D₂ photostability through the formation of cocrystals	
<i>Emma Bello, Eirini Tarelli, Rafael Barbas, Rafel Prohens, Clara Ràfols, Elisabet Fuguet</i>	50
Design of microfluidic devices for fast determination of octanol/water partition coefficients by HPLC	
<i>Elisabet Fuguet, Abdulkarim Albishiri, Joan M. Cabot, Martí Rosés</i>	51

Drug-excipient interactions: evaluation of the binding constants <i>Clara Râfols, Rebeca Ruiz, Elisabet Fuguet</i>	52
Chamelogk: a new experimental-chromatographic chameleonicity descriptor <i>Maura Vallaro, Diego Garcia Jimenez, Matteo Rossi Sebastiano, Giulia Aprato,</i>	53
EPR imaging: A convenient method for <i>in vivo</i> monitoring the efficacy of anti-inflammatory drugs <i>Marko Jovanović, Hadi Waisi, Stevan Blagojević, Vladimir Tajković, Marko Daković, Đura Nakarada, Miloš Mojović</i>	54
Green synthesis of Ag-Fe nanoparticles using <i>cotinus coggygia</i>: Preparation, characterization and photothermal potential <i>Tsenka Grancharova, Stanislava Simeonova, Bissera Pilicheva, Plamen Zagorchev</i>	55
Quantitative measurement of concentration and diffusion properties of molecules using fluorescence correlation spectroscopy <i>J. Z. Jelić, A. Denčevski, M. D. Rabasović, A. J. Krmpot</i>	56
Evaluation of dose dependent oral drug absorption by μFlux <i>Shiori Ishida, Samuel Lee, Karl Box, Balint Sinko, Kiyohiko Sugano</i>	57
Effect of bicarbonate buffer on the precipitation behavior of drugs <i>Hibiki Yamamoto, Kiyohiko Sugano</i>	58
Combined in silico approach in screening of flavonoids database for potential NLRP3 inhibitors <i>Kristina Stevanović, Draginja Radošević, Vladimir Perović, Sanja Glišić</i>	59
Searching for the best way to incorporate the proprietary compound GL-II -73 into the nanoemulsion carrier for prospective parenteral application <i>Jelena Đoković, Bojan Marković, Dishary Sharmin, James M Cook, Miroslav Savić, Snežana Savić</i>	60
Design of tailor-made biocompatible nanocarrier for novel pyrazoloquinolinone ligand (CW-02-79) based on comprehensive evaluation of critical physicochemical descriptors <i>Tijana Stanković, Tanja Ilić, Ivana Pantelić, Anđela Tošić, Jelena Mitrović, James M. Cook, Miroslav Savić, Snežana Savić</i>	61
<i>In silico</i> prediction of liquid-liquid phase separation concentration using abraham solute descriptors <i>Taiga Uekusa, Kiyohiko Sugano</i>	62
Polymorphic conversion of a hydrophobic drug in water: Posaconazole suspensions <i>Matteo Guidetti; Fritz Blatter; Rolf Hilfiker; Annette Bauer-Brandl; Martin Kuentz</i>	63
Challenges in peptide method development Interactions between peptides and HPLC vials <i>Ksenija Žibert, Mateja Filković, Iva Filipović</i>	64

Determination of the nifedipine solubility at biorelevant temperature <i>Ekin Guler, Kaan Altun, Emelhan Rahimoglu, Bolgacem Aybar</i>	65
Clofazimine acid-base solubilization: influence of small organic acids' concentration <i>Igor A. Topalović, Olivera S. Marković, Miloš P. Pešić, Mufaddal Kathawala, Abu T. M. Serajuddin, Alex Avdeef, Tatjana Ž. Verbić</i>	66
¹H-NMR-based serum metabolomics of bipolar disorder patients <i>Nataša Avramović, Katarina Simić, Zoran Miladinović, Nina Todorović, Snežana Trifunović, Aleksandra Gavrilović, Silvana Jovanović, Dejan Gođevac, Ljubodrag Vujisić, Vele Tešević, Ljubica Tasić, Boris Mandić</i>	67
¹H-NMR metabonomic view on schizophrenia <i>Nataša Avramović, Katarina Simić, Zoran Miladinović, Nina Todorović, Snežana Trifunović, Aleksandra Gavrilović, Silvana Jovanović, Dejan Gođevac, Ljubodrag Vujisić, Vele Tešević, Ljubica Tasić, Boris Mandić</i>	68
Salt dissolution, supersaturation, and precipitation kinetics: A comparison between the USP II and the μDISS Profiler™ <i>Laurin Zöllner, Sara Carlert, Eva Karlsson, Anders Borde, Christoph Saal, Jennifer Dressman</i>	69
Development of ophthalmic nanoemulsion for advanced delivery of poorly water- and oil-soluble loteprednol etabonate <i>Josip Ljubica, Bisera Jurišić Dukovski, Drago Špoljarić, Iva Krtalić, Ivan Sušan, Anita Hafner, Ivan Pepić, Jasmina Lovrić</i>	70
Preparation and evaluation of poly-ϵ-caprolactone nanoparticles as carriers for nose-to-brain delivery of lidebenone <i>R. Boyuklieva, B. Pilicheva</i>	71
Directed protein evolution for the investigation of endogenous proteins in synaptic organization and transmission <i>Ivana Trivunovic, Charlotte Rimbault, Ellyn Renou, Cameron Mackereth, Daniel Choquet, Matthieu Sainlos</i>	72
NMR study of haloperidol in weak acid solutions <i>Mladen M. Đurđević, Miloš P. Pešić, Tatjana Ž. Verbić</i>	69

O 20

Navigating towards improved cytotoxicity assessment in nanomedicine development: Shifting from colorimetric to fluorescence-based assays

Ines Nikolić^{1,2}, Marija Petrović¹, Jelena Mitović², Emmanuelle Sublet¹, Olivier Jordan¹, Snežana Savić², Gerrit Borchard¹

¹University of Geneva, Faculty of Science, Section of Pharmaceutical Sciences, Switzerland

²University of Belgrade – Faculty of Pharmacy, Serbia

It is well known that the characterization of nanomedicines can pursue different levels of complexity, both in the development stage and in the quality control process [1]. In line with physicochemical aspects, even more obstacles are encountered in biological safety assessment, while anticipation of their immunogenic potential represents an additional challenge. Moreover, interactions between the test reagents and the nanomaterial have been identified as one of the most important issues in toxicity testing that influence market authorization of nanomedicines, which ought to be resolved [1]. The European Nanomedicine Characterization Laboratory – the reference laboratory for nanomedicines, provides protocols for 2 colorimetric cytotoxicity assays employing LLC-PK1 (porcine kidney epithelial cells) and Hep-G2 (human hepatocarcinoma cells) cell lines. However, the latest recommendations in the field underline the demand for enhancing the testing procedures, while proposing incorporation of immune cells as target cell lines for toxicity evaluation, aiming to provide more reliable conclusions on nanomedicine safety in preclinical level.

In this study 2 inherently different types of pharmaceutical nanosystems were selected: nanoemulsion (NE) and solid lipid nanoparticles (LNP) and subjected to a set of orthogonal toxicity evaluation assays. Adjusted WST-1 (assessing mitochondrial activity as an indicator of cellular well-being) and LDH (lactate dehydrogenase release evaluation as an indicator of cell membrane damage) assays have been performed as the colorimetric tests, while propidium-iodide (PI)-based assay was developed as a fluorescent counterpart (able to directly distinguish live from dead cells), using RAW 246.7 cell line (murine macrophages – immune system cell line). Starting concentration of the tested nanoformulations was 50 % v/v, and they were subsequently diluted with the factor of 2, to create a total of 8 concentrations. Incubation time was 4 h.

Presented assays rely on completely different biological bases. Therefore, their careful combination can address some shortcomings in the *in vitro* evaluations established so far. Although similar toxicity trends were observed regardless the assay used, it was evident that the LDH assay required specific consideration. Since the supernatant is the subject of the analysis (not the cells directly), containing not only the enzyme of interest, but also the nanoformulations, in the wells corresponding to the 3 highest concentration of the NE/LNP pronounced scattering effects were observed. Such an event could be easily overlooked, potentially affecting the conclusions. However, it was overcome by careful design of control and blank wells (each test concentration was coupled with its own blank well containing no cells, but the same concentration of the NE/LNP in the culture medium).

In contrast, absorbance measurements in WST-1 assay were performed in the absence of the NE/LNP, avoiding any interactions or scattering effects. Finally, developed PI-based assay proved to be the most relevant method. Based on the penetration of PI into the dead cell only, attaching to their DNA, the concentration of the dead vs. live cells could be directly estimated. What is more, after the incubation time, the measurements can be performed in the nanoformulation-free environment, surpassing the potential interactions. Notably, cell viability obtained in the PI-based assay followed the same trend as in the WST-1 assay but with significant difference in the obtained values for the first 3 concentrations (Figure 1).




Figure 1. Cytotoxicity assay results for the NE: WST-1 and PI-based assay (*p<0.05)

[1] Simon, 2023. *J Control Rel*, 354, 120-127.

[2] Bremer et al., 2016. Publications Office of the European Union.

Acknowledgement: This research has been supported by the Frankfurt Foundation Quality of Medicines through the project *Quality Requirements for Nanomedicines: Filling the gaps* (Project No. 2023_2) and by the Serbian Science Fund through the project *NanoCellEмоCог* (Grant No. 7749108).